arXiv:1001.2190v2 [cs.IT] 19 May 2010

Characterizations of generalized entropy functions by functional equations

Shigeru Furuichi[1]*
[1]Department of Computer Science and System Analysis,
College of Humanities and Sciences, Nihon University,
3-25-40, Sakurajyousui, Setagaya-ku, Tokyo, 156-8550, Japan

Abstract. We shall show that a two-parameter extended entropy function is characterized by a functional equation. As a corollary of this result, we obtain that the Tsallis entropy function is characterized by a functional equation, which is a different form used in [1] i.e., in Proposition 2.1 in the present paper. We also give an interpretation of the functional equation giving the Tsallis entropy function, in the relation with two non-additive properties.

Keywords : Entropy function, Tsallis entropy, two-parameter extended entropy, characterization and functional equation

2000 Mathematics Subject Classification : 94A17

1 Introduction

Recently, generalized entropies have been studied from the mathematical point of view. The typical generalizations of Shannon entropy [2] are Rényi entropy [3] and Tsallis entropy [4]. The Rényi entropy and the Tsallis entropy are defined by

$$R_q(X) = \frac{1}{1-q}\log\sum_{j=1}^{n} x_j^q, \quad (q \neq 1, q > 0)$$

and

$$S_q(X) = \sum_{j=1}^{n} \frac{x_j - x_j^q}{q-1}, \quad (q \neq 1, q > 0)$$

for a given information source $X = \{x_1, \cdots, x_n\}$ with the probability $p_j \equiv Pr(X = x_j)$. Both entropies recover Shannon entropy:

$$S_1(X) \equiv -\sum_{j=1}^{n} p_j \log p_j$$

in the limit $q \to 1$. The uniqueness theorem for the Tsallis entropy was firstly given in [5] and improved in [6].

Throughout this paper, a parametric extended entropy such as Rényi entropy and Tsallis entropy and so on, is called by a generalized entropy.

---

*E-mail:furuichi@chs.nihon-u.ac.jp

We note that the Rényi entropy has the additivity:

$$R_q(X \times Y) = R_q(X) + R_q(Y)$$

but the Tsallis entropy has the non-additivity:

$$S_q(X \times Y) = S_q(X) + S_q(Y) + (1-q)S_q(X)S_q(Y), \tag{1}$$

where $X \times Y$ means that X and Y are independent random variables. Therefore we have a definitive difference for these entropies, although we have the simple relation between them:

$$R_q(X) = \frac{1}{1-q}\log\left\{1 + (1-q)S_q(X)\right\}, \quad (q \neq 1).$$

The Tsallis entropy is rewritten by

$$S_q(X) = -\sum_{j=1}^{n} p_j^q \ln_q p_j \tag{2}$$

where the q-logarithmic function is defined by

$$\ln_q x \equiv \frac{x^{1-q}-1}{1-q}, \quad (q \neq 1),$$

which uniformly converges to $\log x$ in the limit $q \to 1$.

Since Shannon entropy can be regarded as the expectation value for each value $-\log p_j$, we may consider that the Tsallis entropy can be regarded as the q-expectation value for each value $-\ln_q p_j$, as an anology to Shannon entropy. Where q-expectation value E_q is defined by

$$E_q(X) \equiv \sum_{j=1}^{n} p_j^q x_j.$$

However, the q-expectation value E_q lacks the fundamental property such as $E(1) = 1$, so that it was considered to be inadequate to adopt as a generalized definition of the usual expectation value. Then the noremalized q-expectation value was introduced:

$$E_q^{(nor)}(X) \equiv \frac{\sum_{j=1}^{n} p_j^q x_j}{\sum_{i=1}^{n} p_i^q}$$

and by using this, the normalized Tsallis entropy was defined by

$$S_q^{(nor)}(X) \equiv \frac{S_q(X)}{\sum_{j=1}^{n} p_j^q} = -\frac{\sum_{j=1}^{n} p_j^q \ln_q p_j}{\sum_{i=1}^{n} p_i^q}, \quad (q \neq 1).$$

We easily find that we have the following non-additivity relation for the normalized Tsallis entropy:

$$S_q^{(nor)}(X \times Y) = S_q^{(nor)}(X) + S_q^{(nor)}(Y) + (q-1)S_q^{(nor)}(X)S_q^{(nor)}(Y). \tag{3}$$

As for the details on the mathematical properties of the normalized Tsallis entropy, see [7] for example. The difference between two non-additivities Eq.(1) and Eq.(3) is the signature of the coefficient $1-q$ in the third term of the right hand sides.

We note that the Tsallis entropy is also rewritten by

$$S_q(X) = \sum_{j=1}^{n} p_j \ln_q \frac{1}{p_j} \tag{4}$$

so that we may regard it as the expectation value such as $S_q(X) = E_1\left[\ln_q \frac{1}{p_j}\right]$, where E_1 means the usual expectation value. However, if we adopt this formulation in the definition of the Tsallis conditional entropy, we do not have an important property such as a chain rule. (See [8] for details.) Therefore we often adopt the formulation using the q-expectation value.

As a further generalization, for two positive numbers α and β, a two-parameter extended entropy:

$$S_{\alpha,\beta}(X) \equiv \sum_{j=1}^{n} \frac{x_j^{\alpha} - x_j^{\beta}}{\beta - \alpha}, \quad (\alpha \neq \beta)$$

has been studied in many literatures [9, 10, 11, 12, 13, 14, 15].

In the paper [1], a characterization of the Tsallis entropy function was proven by using the functional equation. In this paper, we shall show that the two-parameter extended entropy function is characterized by the functional equation.

2 A review of the characterization of Tsallis entropy function by the functional equation

The following proposition was originally given in [1] by the simple and elegant proof. Here we give the alternative proof along to the proof given in [16].

Proposition 2.1 ([1]) *If the differentiable nonnegative function f_q with positive parameter $q \in \mathbb{R}$ satisfies the following functional equation:*

$$f_q(xy) + f_q((1-x)y) - f_q(y) = (f_q(x) + f_q(1-x))\, y^q, \qquad (0 < x < 1, \quad 0 < y \leq 1) \tag{5}$$

then the function f_q is uniquely given by

$$f_q(x) = -c_q x^q \ln_q x,$$

where c_q is a nonnegative constant depending only on the parameter q.

Proof:

If we put $y = 1$ in Eq.(5), then we have $f_q(1) = 0$. From here we assume $y \neq 1$. We also put $g_q(t) \equiv \frac{f_q(t)}{t}$. Then we have

$$x g_q(xy) + (1-x) g_q((1-x)y) - g_q(y) = (x g_q(x) + (1-x) g_q(1-x))\, y^{q-1} \tag{6}$$

Putting $x = \frac{1}{2}$ in (6), we have

$$g_q\left(\frac{y}{2}\right) = g_q\left(\frac{1}{2}\right) y^{q-1} + g_q(y).$$

Substituting $\frac{y}{2}$ into y, we have

$$g_q\left(\frac{y}{2^2}\right) = g_q\left(\frac{1}{2}\right)\left(y^{q-1} + \left(\frac{y}{2}\right)^{q-1}\right) + g_q(y).$$

By repeating similar substitutions, we have

$$\begin{aligned} g_q\left(\frac{y}{2^N}\right) &= g_q\left(\frac{1}{2}\right) y^{q-1}\left(1 + \left(\frac{1}{2}\right)^{q-1} + \left(\frac{1}{2}\right)^{2(q-1)} + \cdots + \left(\frac{1}{2}\right)^{(N-1)(q-1)}\right) + g_q(y) \\ &= g_q\left(\frac{1}{2}\right) y^{q-1}\left(\frac{2^{N(1-q)-1}}{2^{1-q} - 1}\right) + g_q(y). \end{aligned}$$

Then we have

$$\lim_{N\to\infty} \frac{g_q\left(\frac{y}{2^N}\right)}{2^N} = 0 \tag{7}$$

due to $q > 0$. Differentiating Eq.(6) by y, we have

$$x^2 g_q(xy) + (1-x)^2 g_q((1-x)y) - g_q'(y) = (q-1)\left(x g_q(x) + (1-x) g_q(1-x)\right) y^{q-2}$$

Putting $y = 1$ in the above equation, we have

$$x^2 g_q'(x) + (1-x)^2 g_q'(1-x) + (1-q)\left(x g_q(x) + (1-x) g_q(1-x)\right) = -c_q, \tag{8}$$

where $c_q = -g_q'(1)$.

By integrating the equation (6) from 2^{-N} to 1 with respect to y and performing the conversion of the variables, we have

$$\int_{2^{-N}x}^{x} g_q(t)dt + \int_{2^{-N}(1-x)}^{1-x} g_q(t)dt - \int_{2^{-N}}^{1} g_q(t)dt = \left(x g_q(x) + (1-x) g_q(1-x)\right) \frac{1-2^{-qN}}{q}. \tag{9}$$

By differentiating the above equation with respect to x, we have

$$\begin{aligned} & g_q(x) - 2^{-N} g_q(2^{-N}x) - g_q(1-x) + 2^{-N} g_q(2^{-N}(1-x)) \\ & = \frac{1-2^{-qN}}{q} \left(g_q(x) + x g_q'(x) - g_q(1-x) - (1-x) g_q'(1-x) \right). \end{aligned}$$

Taking the limit $N \to \infty$ in the above, we have

$$(1-x) g_q'(x) + (1-q) g_q(1-x) = x g_q'(x) + (1-q) g_q(x) \tag{10}$$

thanks to (7). From the equations (8) and (10), we have the following differential equation:

$$x g_q'(x) + (1-q) g_q(x) = -c_q.$$

This differential equation has the following general solution:

$$g_q(x) = -\frac{c_q}{1-q} + d_q x^{q-1},$$

where d_q is a integral constant depending on q. From $g_q(1) = 0$, we have $d_q = \frac{c_q}{1-q}$. Thus we have

$$g_q(x) = c_q \frac{x^{q-1} - 1}{1-q}.$$

Finally we have

$$f_q(x) = c_q \frac{x^q - x}{1-q} = -c_q x^q \ln_q x.$$

From $f_q(x) \geq 0$, we have $c_q \geq 0$.

$\blacksquare$

If we take the limit as $q \to 1$ in Theorem 2.1, we have the following corollary.

Corollary 2.2 ([16]) *If the differentiable nonnegative function f satisfies the following functional equation:*

$$f(xy) + f((1-x)y) - f(y) = \left(f(x) + f(1-x)\right) y, \qquad (0 < x < 1, \quad 0 < y \leq 1) \tag{11}$$

then the function f is uniquely given by

$$f(x) = -cx \log x,$$

where c is a nonnegative constant.

3 Main results

In this section, we give a characterization of a two-parameter extended entropy function by the functional equation. Before we give our main theorem, we review the following result given by Pl.Kannappan [17, 18].

Proposition 3.1 ([17, 18]) *Let two probability distributions* $(p_1, \cdots, p_n)$ *and* $(q_1, \cdots, q_m)$*. If the measureable function* $f : (0,1) \to \mathbb{R}$ *satisfies*

$$\sum_{i=1}^{n}\sum_{j=1}^{m} f(p_i q_j) = \sum_{i=1}^{n} p_i^{\alpha} \sum_{j=1}^{m} f(q_j) + \sum_{j=1}^{m} q_j^{\beta} \sum_{i=1}^{n} f(p_i), \tag{12}$$

for all $(p_1, \cdots, p_n)$ *and* $(q_1, \cdots, q_m)$*, then the function* f *is given by*

$$f(p) = \begin{cases} c\left(p^{\alpha} - p^{\beta}\right), & \alpha \neq \beta, \\ cp^{\alpha}\log p, & \alpha = \beta, \\ cp\log p + b\left(mn - m - n\right)p + b, & \alpha = \beta = 1. \end{cases}$$

where c *and* b *are arbitrary constants.*

In the following theorem, we adopt a simpler condition than Eq.(12).

Theorem 3.2 *If the differentiable nonnegative function* $f_{\alpha,\beta}$ *with two positive parameters* $\alpha, \beta \in \mathbb{R}$ *satisfies the following functional equation:*

$$f_{\alpha,\beta}(xy) = x^{\alpha} f_{\alpha,\beta}(y) + y^{\beta} f_{\alpha,\beta}(x), \qquad (0 < x, y \leq 1) \tag{13}$$

then the function $f_{\alpha,\beta}$ *is uniquely given by*

$$f_{\alpha,\beta}(x) = c_{\alpha,\beta}\frac{x^{\beta} - x^{\alpha}}{\alpha - \beta}, \quad (\alpha \neq \beta)$$

and

$$f_{\alpha}(x) = -c_{\alpha}x^{\alpha}\log x, \quad (\alpha = \beta)$$

where $c_{\alpha,\beta}$ *and* c_{α} *are nonnegative constants depending only on the parameters* α *(and* β*).*

Proof:

If we put $y = 1$, then we have $f_{\alpha,\beta}(1) = 0$ due to $x > 0$. By differentiating Eq.(13) with respect to y, we have

$$x f'_{\alpha,\beta}(xy) = x^{\alpha} f'_{\alpha,\beta}(y) + \beta y^{\beta-1} f_{\alpha,\beta}(x) \tag{14}$$

Putting $y = 1$ in Eq.(14), we have the following differential equation:

$$x f'_{\alpha,\beta}(x) - \beta f_{\alpha,\beta}(x) = -c_{\alpha,\beta}x^{\alpha}, \tag{15}$$

where we put $c_{\alpha,\beta} \equiv -f'_{\alpha,\beta}(1)$. Eq.(15) can be deformed as follows.

$$x^{\beta+1}\left(x^{-\beta} f_{\alpha,\beta}(x)\right)' = -c_{\alpha,\beta}x^{\alpha}.$$

That is, we have,

$$\left(x^{-\beta} f_{\alpha,\beta}(x)\right)' = -c_{\alpha,\beta}x^{\alpha-\beta-1}.$$

Integrating both sides on the above equation with respect to x, we have

$$x^{-\beta} f_{\alpha,\beta}(x) = -\frac{c_{\alpha,\beta}}{\alpha-\beta} x^{\alpha-\beta} + d_{\alpha,\beta},$$

where $d_{\alpha,\beta}$ is a integral constant depending on α and β. Therefore we have

$$f_{\alpha,\beta}(x) = -\frac{c_{\alpha,\beta}}{\alpha-\beta} x^{\alpha} + d_{\alpha,\beta} x^{\beta},$$

By $f_{\alpha,\beta}(1) = 0$, we have $d_{\alpha,\beta} = \frac{c_{\alpha,\beta}}{\alpha-\beta}$. Thus we have

$$f_{\alpha,\beta}(x) = \frac{c_{\alpha,\beta}}{\alpha-\beta}\left(x^{\beta} - x^{\alpha}\right).$$

Also by $f_{\alpha,\beta}(x) \geq 0$, we have $c_{\alpha,\beta} \geq 0$.

As for the case of $\alpha = \beta$, we can prove by the similar way.

■

If we take $\alpha = q, \beta = 1$ or $\alpha = 1, \beta = q$ in Theorem 3.2, we have the following corollary.

Corollary 3.3 *If the differentiable nonnegative function f_q with a positive parameter $q \in \mathbb{R}$ satisfies the following functional equation:*

$$f_q(xy) = x^q f_q(y) + y f_q(x), \qquad (0 < x, y \leq 1, \quad q \neq 1) \tag{16}$$

then the function f_q is uniquely given by

$$f_q(x) = -c_q x^q \ln_q x$$

where c_q is a nonnegative constant depending only on the parameter q.

Here we give an interpretation of the functional equation (16) from the view of Tsallis statistics.

Remark 3.4 *Replacing x by y each other in Eq.(16) and summing two functional equations, we have*

$$f_q(xy) = \left(\frac{x^q + x}{2}\right) f_q(y) + \left(\frac{y^q + y}{2}\right) f_q(x), \qquad (0 < x, y \leq 1, \quad q \neq 1). \tag{17}$$

Then the functional equation (17) can be regarded as the sum of two following functional equations:

$$f_q(xy) = y f_q(x) + x f_q(y) + (1-q) f_q(x) f_q(y) \tag{18}$$

$$f_q(xy) = y^q f_q(x) + x^q f_q(y) + (q-1) f_q(x) f_q(y) \tag{19}$$

Two equations (18) and (19) imply the following equations for $i = 1, \cdots, n$ and $j = 1, \cdots, m$.

$$f_q(x_i y_j) = y_j f_q(x_i) + x_i f_q(y_j) + (1-q) f_q(x_i) f_q(y_j) \tag{20}$$

$$f_q(x_i y_j) = y_j^q f_q(x_i) + x_i^q f_q(y_j) + (q-1) f_q(x_i) f_q(y_j) \tag{21}$$

Taking the sum for Eq.(20) and Eq.(21) on i and j with simple calculations, we have two non-additivity relations given in Eq.(1) and Eq.(3), where we put $c_q = 1$ for a simplicity. Therefore we can conclude that two functional equations (18) and (19), which are the essential parts of the non-additivity relations Eq.(1) and Eq.(3), characterize the Tsallis entropy function. In other words, the Tsallis entropy function can be characterized by two non-additivity relations Eq.(1) and Eq.(3).

If we again take the limit as $q \to 1$ in Corollary 3.3, we have the following corollary.

Corollary 3.5 *If the differentiable nonnegative function f satisfies the following functional equation:*

$$f(xy) = yf(x) + xf(y), \qquad (0 < x, y \leq 1) \tag{22}$$

then the function f is uniquely given by

$$f(x) = -cx \log x$$

where c is a nonnegative constant.

Ackowledgement

The author was partially supported by the Japanese Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Encouragement of Young Scientists (B) 20740067.

References

[1] H.Suyari and M.Tsukada, Tsallis differential entropy and divergences derived from the generalized shannon-Khinchin axioms, Proc.2009,IEEE-ISIT,pp.149-153.

[2] C.E.Shannon, A mathematical theory of communication, Bell Syst.Tech.J.,Vol.27(1948), pp.379-423 and pp.623-656.

[3] A.Rényi, On measures of entropy and information, in Proc. 4th Berkeley Symp., Mathematical and Statistical Probability, Berkeley, CA: Univ. Calif. Press, vol. 1(1961), pp. 547-561.

[4] C. Tsallis, Possible generalization of Bolzmann-Gibbs statistics, J.Stat. Phys., Vol. 52(1988), pp. 479-487.

[5] H.Suyari, Generalization of Shannon-Khinchin axioms to nonextensive systems and the uniqueness theorem for the nonextensive entropy, IEEE Trans. Information Theory, Vol.50(2004), pp.1783-1787.

[6] S.Furuichi, On uniqueness theorems for Tsallis entropy and Tsallis relative entropy, IEEE Trans. Information Theory, Vol.51(2005), pp.3638-3645.

[7] H.Suyari, Nonextensive entropies derived from form invariance of pseudoadditivity, Phys.Rev.E.,Vol.65(2002), 066118.

[8] S.Furuichi, Information theoretical properties of Tsallis entropies, J.Math.Phys., Vol.47(2006), 023302.

[9] B.D.Sharma and L.J.Taneja,Entropy of type (α, β) and other generalized measures in information theory, Metrika, Vol.22(1975),pp.205-215.

[10] D.P.Mittal, On some functional equations concerning entropy, directed divergence and inaccuracy, Metrika, Vol.22(1975),pp.35-45.

[11] E.P.Borges and I.Roditi,A family of nonextensive entropies, Phys.Lett.A, Vol.246(1998),pp.399-402.

[12] G.Kaniadakis, M.Lissia and A.M.Scarfone, Deformed logarithms and entropies, Physica A, Vol.340(2004),pp.41-49.

[13] G.Kaniadakis, M.Lissia and A.M.Scarfone, Two-parameter deformations of logarithm, exponential, and entropy: A consistent framework for generalized statistical mechanics, Phys.Rev.E, Vol.71(2005),046128.

[14] T.Wada and H.Suyari, A two-parameter generalization of Shannon-Khinchin axioms and the uniqueness theorem, Phys.Lett.A, Vol.368(2007),pp.199-205.

[15] S.Furuichi, An axiomatic characterization of a two-parameter extended relative entropy, arXiv:0810.5399v2.

[16] Y. Horibe, Entropy of terminal distributions and the Fibonacci trees, Fibonacci Quart. Vol.26 (1988), pp.135-140.

[17] Pl. Kannappan, An application of a differential equation in information theory, Gla.Mat.,Vol.14(1979),pp.269-274.

[18] Pl.Kannappan, Functional equations and inequalities with applications, Springer, 2009.